




Why Rodent Control Matters

Rats and mice aren't just a nuisance – they're a serious threat to health, safety and productivity.

Rodents can:

- Damage buildings and equipment
- Consume and contaminate stored food
- Spread harmful bacteria, viruses, parasites and disease

Careful, precise application of rodenticides improves control and protects people, animals, and the environment.

What to Do

Know your target rodent species

Always follow label instructions

Wear PPE

Use bait stations to help protect children, pets, and non-target species

Store rodenticides safely

What to Avoid

Applying more rodenticide than needed

Placing bait where pets or children could access it

Ignoring signs of exposure to the user

Using rodenticides near food preparation areas

Racumin® Foam

Our ready-to-use, contact rodenticide that targets rats and mice through grooming behaviour—ideal for hard-to-reach areas where traditional baits may be less effective.

✓ **Do's**

- Apply in known rat runways, passages, entrance points / holes, covered tracks and wall cavities
- Ensure rodents continue using entrance points by not sealing holes with foam
- Apply at least 2m from food storage or preparation areas
- Protect from sunlight

✗ **Dont's**

- Not for areas frequently cleaned with water, or sensitive surfaces that can stain
- Don't expose to temperatures below 5°C or above 50°C
- Never spray on a naked flame or any incandescent material
- Do not use on unprotected electrical cables

Harmonix® Rodent Paste

This fast-acting, non-anticoagulant bait features the stop-feeding effect. It's suitable for use indoors, outdoors, and in burrows, making it ideal for IPM programs.

✓ **Do's**

- Use in inactive areas to help track rodent activity
- Evaluate risks before application
- Ensure adequate bait for a lethal dose before the stop-feeding effect begins
- Monitor bait uptake. Check every 1-2 days, adding bait every 7 days until consumption stops

✗ **Dont's**

- Never apply without a risk assessment first
- Do not underdose: Insufficient bait may trigger the stop-feeding effect before a lethal dose is consumed
- Don't assume one-time baiting is enough: Ongoing monitoring and replenishment are key

Racumin® Expert

This multi-feed anticoagulant bait is highly palatable. Its delayed action is ideal for cautious rodents, especially in low to moderate infestations.

✓ **Do's**

- Identify the species and its activity zones first
- Place bait boxes in the most effective locations
- Adapt baiting strategy if needed
- Monitor bait consumption; every 2-3 days for voles, and every 5-7 days for rats.

✗ **Dont's**

- Don't ignore the condition of bait. It's less effective after 7-15 days
- Never treat with old bait. Use new product every 2-3 months

Rodilon® Block

This highly palatable, single-feed rodenticide has no known resistance and comes in a durable, moisture-resistant formulation.

✓ **Do's**

- Remove alternative food and water sources
- Place where there's signs of presence (e.g. droppings, greasy spots, hair, and footprints)
- Maintain permanent priming points

✗ **Dont's**

- Don't ignore the condition of bait. It's less effective after 7-15 days
- Never treat with old bait. Use new product every 2-3 months

Rodilon® Soft Blocks

Our single-feed, highly palatable bait is a soft paste that's ideal for indoor and perimeter control, and dry environments.

✓ **Do's**

- Expect visible effects 3-5 days after rodents first consume the bait
- Be patient. With Neophobia, it may take a few days for rats to feed
- Place enough bait for a lethal dose in a single feed

✗ **Dont's**

- Don't assume immediate results. Delayed action is normal
- Do not remove bait too soon. Give rodents time to discover and consume it

Rodilon® Pellets

These ready-to-use, moisture-resistant pellets offer high palatability and durability. The grain-based formulation is ideal for use in hidden areas and transit zones.

✓ **Do's**

- Use the correct quantities for rats and mice
- Use dedicated tools for handling
- Adjust dosage based on the infestation level
- Only use in tamper-resistant bait stations
- Maintain bait availability, monitor and replace as needed

✗ **Dont's**

- Never reuse empty containers or utensils.
- Bait or carcasses should not be left unattended.

Racumin® Powder

This multiple-dose anticoagulant is a concentrated ready-to-use powder that adheres to fur and is ingested during grooming. It's ideal for cautious rats and mice in hard-to-bait areas.

✓ **Do's**

- Expect visible effects 3-5 days after rodents first consume the bait
- Be patient. With Neophobia, it may take a few days for rats to feed
- Place enough bait for a lethal dose in a single feed

✗ **Dont's**

- Don't assume immediate results. Delayed action is normal
- Do not remove bait too soon. Give rodents time to discover and consume it

For more information, please visit www.uk.envu.com

Envu and Rodilon® and Racumin® and Harmonix® are all registered trademarks of Environmental Science US LLC or one of its affiliates. Always read the label and product information before use. For further information, including contact details, visit www.envu.com.

© 2025 Environmental Science U.S. LLC.

envu™