

2020-3386
2020-10-23



GROUP	4	INSECTICIDE
-------	---	-------------

QuickBayt® Granular Fly Bait

Insecticide

AGRICULTURAL

FOR INDOOR USE AT LIVESTOCK FACILITIES ONLY

Attracts and Kills House Flies

GUARANTEE:

Imidacloprid.0.5%
Muscalure...0.09%

REGISTRATION NUMBER 32234 PEST CONTROL PRODUCTS ACT

READ THE LABEL BEFORE USING

KEEP OUT OF REACH OF CHILDREN

NET CONTENTS: 2.26 kg

Bayer CropScience Inc.
Suite 200, 160 Quarry Park Blvd. S.E.
Calgary, Alberta T2C 3G3

Product Information: 1-888-283-6847
www.es.bayer.ca

In case of spills, poisoning or fire, telephone emergency response number
1-800-334-7577 (24 hours a day).

GENERAL INFORMATION

Attracts and kills house flies inside livestock facilities

Ready-to-use bait (containing 0.5% imidacloprid)

Contains a specially formulated mixture of attractants

Contains the attractant Muscalure

Contains Bitrex® bittering agent to help prevent ingestion by animals, children, and pets.


DIRECTIONS FOR USE

Use QuickBayt Granular Fly Bait as part of an overall integrated pest management program for house flies which may include:

- Proper sanitation and manure management.
- Environmental controls, such as proper light management and exclusion.
- Other registered pest control products.

DO NOT formulate this product into other end-use products.

TO ATTRACT AND KILL HOUSE FLIES	
WHERE TO APPLY	<p>FOR INDOOR USE ONLY.</p> <p>For use inside livestock facilities such as broiler houses, feedlots, livestock housing structures, and horse stables.</p> <p>DO NOT place bait in areas accessible to livestock, pets and children.</p> <p>Apply QuickBayt Granular Fly Bait in locations where house flies congregate. In caged layer houses, apply only in the walkways.</p> <p>Avoid application in cool locations because house flies prefer warm sites for resting. Avoid application in areas with air movement such as near fans.</p> <p>When used in areas accessible to birds, bait must be placed in a tamper- resistant bait station that prevents birds from consuming the bait.</p>
WHEN TO APPLY	<p>Time applications to begin at the start of the season before fly populations have reached their peak.</p>
USE RATE	<p>200 g QuickBayt Granular Fly Bait /100 m²</p>
METHODS OF APPLICATION	<p>BROADCAST APPLICATION:</p> <p>Scatter the bait at a rate of 200 g/100m² directly from the container onto dry level surfaces so that the individual granules lie near each other without forming small piles. DO NOT use this method of application in horse stables.</p> <p>BAIT STATIONS:</p> <p>Place in tamper-resistant fly bait station. Add 50 g of bait per station. Use 1 bait station to every 25 m². Secure bait stations at least 1.25 m above the ground. Bait stations should be inaccessible to livestock, children and pets.</p>

	<p>Treated baits must be placed in tamper-resistant bait stations OR locations NOT accessible to children, pets, livestock or wildlife.</p> <p>Tamper-resistant bait stations must:</p> <ol style="list-style-type: none"> 1. be constructed of high-strength material that is hard for children or animals to destroy; 2. be constructed so that the bait cannot be shaken out; 3. have an entrance which prevents children and non-target animals from reaching bait; 4. have an access panel which fastens securely and locks; 5. be able to be securely fastened to a surface; and 6. be clearly labelled: DANGER POISON.
RESTRICTIONS 	<p>Treated baits must be placed in tamper-resistant bait stations OR locations NOT accessible to children, pets, livestock or wildlife.</p> <p>Do not allow bait to come into contact with food or water, livestock feed or their water.</p> <p>Do not apply to milking equipment.</p> <p>Do not apply inside food processing plants, restaurants or other commercial facilities.</p>

RESISTANCE MANAGEMENT RECOMMENDATIONS

For resistance management, please note that QuickBayt Granular Fly Bait contains a Group 4 insecticide. Any insect population may contain individuals naturally resistant to QuickBayt Granular Fly Bait and other Group 4 insecticides. The resistant individuals may dominate the insect population if this group of insecticides is used repeatedly in the same location. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance management strategies should be followed.

To delay insecticide resistance:

- Where possible, rotate the use of QuickBayt Granular Fly Bait or other Group 4 insecticides with different groups that control the same pests.
- Insecticide use should be based on an Integrated Pest Management (IPM) program that includes scouting, record keeping, and considers cultural, biological and other chemical control practices.
- Monitor treated pest populations for resistance development.
- Contact your local extension specialist for any additional pesticide resistance- management and/or IPM recommendations for the specific site and pest problems in your area.

For further information and to report suspected resistance, contact Bayer CropScience Inc. at 1-888-283-6847 or at www.cropscience.bayer.ca.

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN

DO NOT APPLY IN THE PRESENCE OF CHILDREN

DO NOT APPLY TO AREAS ACCESSIBLE TO CHILDREN

Wear a long-sleeved shirt, long pants and chemical resistant gloves during application and clean

up activities.

Avoid contamination of food and feed.

Avoid contact with eyes, skin or clothing.

Wash thoroughly with soap and water after handling.

ENVIRONMENTAL HAZARDS

Toxic to aquatic organisms, birds, and bees. Do not contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

Apply this product only in accordance with this label.

FIRST AID

IF SWALLOWED: Call a poison control centre or doctor immediately for treatment advice.

Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION

No specific antidote is available. Treat the patient symptomatically.

STORAGE

Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store in original container and out of the reach of children, preferably in a locked storage area.

DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

1. Thoroughly empty the contents of the container before disposal.
2. Follow provincial instructions for any required additional cleaning of the container prior to its disposal.
3. Make the empty container unsuitable for further use.
4. Dispose of the container in accordance with provincial requirements.
5. For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

ACCIDENTAL SPILLS

Handle and open container in a manner as to avoid spillage. If the container is leaking or material is spilled for any reason or cause, carefully sweep up spilled bait and wash exposed area. In spill or leak incidents, keep unauthorized people away. You may contact the 24-hour Emergency Phone number for decontamination procedures or any other assistance that may be necessary.

The 24-hour Emergency Phone number is 1-800-334-7577.

NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

® TM see www.bayer.ca/tm-mc

All other trademarks are the property of their respective owners.