



Bayer Plant Protection Calendar

We'll meet you there
with innovative turf solutions.



PUTTING GREEN PROGRAM

Application date	Fungicide and rate	Mode of action	Targets
Mid-April	Dedicate Stressgard (48 mL)	DMI + Qol	Microdochium / Red leaf spot <i>SPRING CLEANUP</i>
Mid-June	Mirage Stressgard (32 mL) Signature XTRA Stressgard (60 – 120 g)	DMI + Phosphonate	Dollar spot / Anthracnose / Patch diseases <i>SUMMER DECLINE</i>
Mid-July	Exteris Stressgard (140 mL) Signature XTRA Stressgard (60 – 120 g)	SDHI + Qol Phosphonate	Dollar spot / Brown patch <i>BROAD SPECTRUM PROTECTION</i>
Early August	Mirage Stressgard (32 mL) Signature XTRA Stressgard (60 – 120 g)	DMI + Phosphonate	Dollar spot / Anthracnose / Patch diseases / <i>SUMMER DECLINE</i>
Late August	Exteris Stressgard (140 mL)	SDHI + Qol	Anthracnose / Dollar spot <i>PLANT HEALTH</i>
Mid-September	Mirage Stressgard (32 mL)	DMI	Anthracnose / Dollar spot <i>FALL PREP</i>
October	Dedicate Stressgard (32-64 mL)	DMI + Qol	Late season dollar spot / Microdochium <i>GO IN CLEAN</i>
November	Dedicate Stressgard (48-64 mL) + Fludioxonil	Qol + DMI + Phenylpyrrole	Typhula / Pink snow moulds <i>WINTER PROTECTION</i>

FAIRWAY PROGRAM

Application date	Fungicide and rate	Mode of action	Targets
June	Mirage Stressgard (32 mL)	DMI	Dollar spot / Anthracnose <i>SUMMER PREP</i>
July	Exteris Stressgard (140 mL)	SDHI + Qol	Dollar spot / Anthracnose / Leaf spot <i>PEAK SEASON DOLLAR SPOT</i>
August	Mirage Stressgard (32 mL)	DMI	Dollar spot / Anthracnose / Brown spot <i>SUMMER CLEAN UP</i>
November	Dedicate Stressgard (48 mL)	DMI + Qol	Typhula / Pink snow moulds <i>WINTER PROTECTION</i>

INSECTICIDE PROGRAM

Application date	Insecticide and rate	Mode of action	Targets
Early June	Tetrino (11.6-23.2 mL)	Diamide	1st generation ABW
July	Tetrino (11.6-23.2 mL)	Diamide	White grubs / 2nd generation ABW / Chinch bugs / Cutworms / Sod webworms
All summer	Maxforce Quantum	Neonicotinoid	Spot treatment for ant mounds

Suggested rates are based on 100 m². Interval of control and suggested programs are based on average historical disease incidence in temperate regions in Canada. Information is believed to be accurate but is not guaranteed. Bayer assumes no responsibility or liability for any action taken as a result of any information or advice contained in this plan, and any action taken is solely at the liability and responsibility of the user.

ALWAYS READ AND FOLLOW PESTICIDE LABEL DIRECTIONS.
Bayer, the Bayer Cross, Banol®, Compass® 50 WG, Exteris™ Stressgard®, Signature XTRA™, Mirage™ Stressgard®, Dedicate™ Stressgard®, Deltagard® SC, Tetrino™, and Maxforce® Quantum is a trademark of Bayer Group. Bayer CropScience Inc. is a member of CropLife Canada. ©2022 Bayer Group. All rights reserved. | 82405-02/22



ALL RATES ARE PER 100 m²

Symptoms Visible Periods of Pathogen Activity

ALL RATES ARE PER 100 m²

Periods of Egg Laying Periods of Feeding

