



Maxforce[®]
PRIME

Contents :
30 g e

INSECTICIDE

✓ A ready to use insecticidal gel bait for the control of cockroaches, nymphs and adults.

Imidacloprid 2.217% (2.15% pure)
UK-2019-1193

Read attached instructions before use.
FOR USE ONLY AS AN INSECTICIDE (PT18). FOR USE ONLY BY PROFESSIONAL OPERATORS.
For indoor use only.

Maxforce Prime is a ready to use insecticidal gel bait containing 2.217% (2.15% pure) Imidacloprid for the control of cockroaches nymphs and adults. Maxforce Prime cartridges are designed for the controlled placement of bait indoors in the following situations:

Indoors

- Industrial / commercial premises
- Households / private areas
- Public (e.g. Hospitals, Nursing Homes)

Maxforce Prime is for the control of German (*Blattella germanica*), Brown-Banded (*Supella longipalpa*), Oriental (*Blatta orientalis*) and American (*Periplaneta americana*) cockroaches.

For 24 hour emergency information contact:
Bayer CropScience Ltd. Telephone: 00800 1020 3333

Approval Holder: Bayer CropScience Limited,
230 Cambridge Science Park,
Milton Road, Cambridge, CB4 0WB.

Further information is available from: Tel: 00800 12149451
www.environmentalscience.bayer.co.uk for Safety Data Sheet & larger label.

Manufacturing date / Batch number: see the cartridge.

Maxforce[®] is a Registered Trade Mark of Bayer. © Copyright of Bayer 2019.



WARNING

GB79225253E - ARTICLE 81756287 - A5

Bayer



Maxforce
PRIME

1

INSECTICIDE

✓ **A ready to use insecticidal gel bait for the control of cockroaches, nymphs and adults**

Contains 2.217% (2.15% pure) Imidacloprid.

4 x 30 g e

Cockroach Control



Safety information
MAXFORCE® PRIME - Contains 2.217% (2.15% pure) Imidacloprid.
May cause an allergic skin reaction.
Very toxic to aquatic life with long lasting effects.
Avoid release into the environment.
Wear protective gloves/ protective clothing/ eye protection/ face protection.
IF ON SKIN: Wash with plenty of soap and water.
If skin irritation or rash occurs: get medical advice/attention.
Specific treatment (see on this label).
Take off contaminated clothing and wash it before reuse.
Collect spillage.
Dispose of contents/container to a licensed hazardous-waste disposal contractor or collection site except for empty clean containers which can be disposed of as non-hazardous waste.
Contains: 1,2-Benzisothiazol-3(2H)-one and Mixture of: 5-chloro-2-methyl-4-isothiazolin-3-one and 2-methyl-4-isothiazolin-3-one.
To avoid risks to human health and the environment, comply with the instructions for use.
UK-2019-1193



WARNING

For 24 hour emergency information contact:
Bayer CropScience Ltd. Telephone: 00800 1020 3333
Approval Holder: Bayer CropScience Limited,
230 Cambridge Science Park, Milton Road, Cambridge, CB4 0WB.
Further information is available from: Tel: 00800 12149451
www.environmentalscience.bayer.co.uk for Safety Data Sheet & larger label.

Manufacturing date / Batch number: see the cartridge.

Maxforce® is a Registered Trade Mark of Bayer. © Copyright of Bayer 2019.

Bayer

GB81773343D - ARTICLE 81756287 - A5

FOR USE ONLY AS AN INSECTICIDE (PT18). FOR USE ONLY BY PROFESSIONAL OPERATORS.

For indoor use only.

Use biocides safely and sustainably.

Always read the label or leaflet before use and follow all the instructions provided.

Maxforce Prime is a ready to use insecticidal gel bait for the control of cockroaches, nymphs and adults.

Maxforce Prime contains 2.217% (2.15% pure) Imidacloprid, along with foodstuffs which are particularly attractive to cockroaches and colloidal materials which maintain attractiveness of the bait and provide cockroaches with a source of liquid.

Maxforce Prime withstands a broad range of temperatures. It is odourless.

Maxforce Prime cartridges are designed for the controlled placement of bait indoors in the following situations:

Indoors

- Industrial / commercial premises
- Households / private areas
- Public (e.g. Hospitals, Nursing Homes)

Maxforce Prime is for use Indoors for the control of:

- German Cockroaches (*Blattella germanica*)
- Brown-Banded Cockroaches (*Supella longipalpa*)
- Oriental Cockroaches (*Blatta orientalis*)
- American Cockroaches (*Periplaneta americana*)

DIRECTIONS FOR USE

For indoor use only.

Apply Maxforce Prime as spots or thin ribbons to surfaces in identified target areas within buildings, it can be placed in corners, cracks and crevices and voids.

The application rate is adjusted to the size of the infestation.

One spot = 0.1 g (approx. 7 mm diameter).

A thin ribbon equating to 1 spot can alternatively be applied.

Application equipment:

Maxforce Prime is dispensed using a suitable gel applicator. When the treatment has been completed remove the empty cartridge and dispose of safely. If the cartridge is only partially used, remove it from the applicator, seal it with the end cap provided and store as directed.

Application Rate:

Cockroach Species	Application rate for given density of infestation * per m ²	
	Low	High
Small e.g. German cockroach, Brown banded cockroach	One spot x 0.1 g	Two spots x 0.1 g
Large e.g. Oriental cockroach, American cockroach	Two spots x 0.1 g	Three spots x 0.1 g

** Maximum application rate of 3 x 0.1g spots per square metre, with a maximum repeat application of 3 – 11 times per year per building.*

Maximum rates of:

- 60 x 0.1 g spots (i.e. 6 g of bait) in a domestic dwelling;
- 270 x 0.1 g spots (i.e. 27.0 g pf bait) in a larger building should not be exceeded.

Should higher amounts of bait be required, then a bait station should be used indoors.

General:

Do not apply directly to surfaces on which food or feed is stored, prepared or eaten.

Use only in positions inaccessible to children and animals.

Prevent access to bait by children and animals.

Wash hands and exposed skin before meals and after use.

Hazardous to bees.

Wear protective chemical resistant gloves during product handling phase, Nitrile gloves.

Do not apply to areas susceptible to routine wet cleaning.

Do not use the product in areas where resistance to the active substance(s) is suspected or established.

In case of re-infestation, reapply the product without exceeding the maximum number of treatments authorised per year.

Inform the registration holder if the treatment is ineffective.

Before treatment, remove all other sources of food for cockroaches (waste, food scraps ...) from the infested area to encourage the ingestion of the gel.

Do not apply the product on absorbent surfaces.

Do not expose bait drops to sunlight or heat (i.e radiator).

Avoid continuous use of the product.

Alternate products containing active substances with a different mode of action.

If the infestation persists despite following the instructions on the label/leaflet, contact a pest control professional.

Use only in concealed areas difficult to access and kept away from water, such as cracks and crevices.

Adopt integrated pest management methods such as the combination of chemical, physical control methods and other public health measures, taking into account local specificities (climatic conditions, target species, conditions of use, etc).

Further Information:

An effect on the cockroach population can be expected very quickly, dead cockroaches may be evident within 24 hours of treatment.

Maximum levels of control are achieved between six days and several months after treatment depending upon the level of infestation and as long as the gel is present.

STORAGE:

Store in original container.

Keep in a cool, dry, well ventilated area.

Protect from frost.

Store away from light.

Shelf life of up to 24 months.

DISPOSAL:

This material and its container must be disposed of in a safe way.

Do not discharge unused product on the ground, into watercourses, into pipes (sink, toilets ...) nor down the drains.

If removing any residual hardened bait by sponging or wet cloth, all bait and used materials (sponge, cloth etc.) must be disposed of safely as domestic waste.

FIRST AID

General:

Move out of dangerous area. Place and transport victim in stable position (lying sideways). Remove contaminated clothing immediately and dispose of safely.

Ingestion:

Wash out mouth with water. Contact poison treatment specialist. Seek medical advice immediately if symptoms occur and/or large quantities have been ingested. In case of impaired consciousness place in recovery position and seek medical advice immediately. Do not give fluids or induce vomiting. Keep container or label available. If necessary take the person to hospital and show the labelling or packaging. Do not leave poisoned person alone.

Skin Contact:

Remove contaminated clothing and shoes. Wash contaminated skin with soap and water. Contact poison treatment specialist if symptoms occur. Seek medical attention if irritation develops and persists.

Eye Contact:

Rinse immediately with plenty of water, also under the eyelids, for at least 15 minutes. Do not forget to remove any contact lenses.



Maxforce[®]
PRIME



INSECTICIDE

✓ **A ready to use insecticidal gel bait for the control of cockroaches, nymphs and adults.**

Contains 2.217% (2.15% pure) Imidacloprid.

Cockroach Control



10 x 4 x 30 g e



Maxforce[®]
PRIME

ARTICLE 81756287

**FOR USE ONLY AS AN INSECTICIDE (PT18). FOR USE ONLY BY PROFESSIONAL OPERATORS.
For indoor use only.**

Safety information

MAXFORCE[®] PRIME

Contains 2.217% (2.15% pure) Imidacloprid.

May cause an allergic skin reaction.

Very toxic to aquatic life with long lasting effects.

Avoid release into the environment.

Wear protective gloves/ protective clothing/ eye protection/ face protection.

IF ON SKIN: Wash with plenty of soap and water.

If skin irritation or rash occurs: get medical advice/attention.

Specific treatment (see on this label).

Take off contaminated clothing and wash it before reuse.

Collect spillage.

Dispose of contents/container to a licensed hazardous-waste disposal contractor or collection site except for empty clean containers which can be disposed of as non-hazardous waste.

Contains: 1,2-Benzisothiazol-3(2H)-one and Mixture of: 5-chloro-2-methyl-4-isothiazolin-3-one and 2-methyl-4-isothiazolin-3-one.

To avoid risks to human health and the environment, comply with the instructions for use.

UK-2019-1193



WARNING

For 24 hour emergency information contact: Bayer CropScience Ltd. Telephone: 00800 1020 3333

Approval Holder: Bayer CropScience Limited,
230 Cambridge Science Park,
Milton Road, Cambridge, CB4 0WB.

Further information is available from: Tel: 00800 12149451
www.environmentalscience.bayer.co.uk for Safety Data Sheet & larger label.

Manufacturing date / Batch number: see the cartridge.

Maxforce[®] is a Registered Trade Mark of Bayer. © Copyright of Bayer 2019.

Bayer

GB80219903E - A5