



GROUP	23	INSECTICIDE
-------	----	-------------

**FIFRA Section 24(c) Special Local Need Label
FOR DISTRIBUTION AND USE ONLY WITHIN FLORIDA**

REVIEWED
By Ashok Shahane at 8:48 pm, Oct 24, 2022

For Control of Bermudagrass Stunt Mites in Golf Courses and Sod Farms

Kontos®

ACTIVE INGREDIENT:

Spirotetramat: *cis*-3-(2,5-dimethylphenyl)-8-methoxy-2-oxo-1-azaspiro[4,5]dec-3-en-4-yl-ethyl carbonate.....22.4%

OTHER INGREDIENTS:.....77.6%
100.0%

Contains 2 pounds of spirotetramat per US gallon, or 240 grams per liter.

EPA Reg. No. 432-1471

SLN. No. FL-220002

CAUTION

THIS LABEL IS VALID UNTIL DECEMBER 31, 2027 UNLESS OTHERWISE AMENDED, WITHDRAWN, CANCELED, OR SUSPENDED.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This label must be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

Pests Controlled	Product Rate (fl oz/A)
Bermudagrass Stunt Mites	Apply up to 5 fl oz of product (150 mL, 0.08 lb active ingredient) per acre in a single application, every two weeks, for a maximum of 25 fl oz of product (750 mL, 0.4 lb active ingredient) per acre per year.
Application Information	
Use properly calibrated application equipment (ground sprayers including hand-held and backpack sprayers) that will produce a uniform coarse droplet spray using a low-pressure setting to eliminate off-target drift.	
Apply in sufficient water to provide optimum distribution in a treated area. Make ground applications in a minimum of 10 gallons of spray solution per acre. Or delivered in a spray volume of 2 gallons per 1,000 sq. ft.	
Start treatments prior to establishment of high pest population and reapply on an as-needed basis. Apply when pests first appear or when damage is first noticed. Spray thoroughly.	
The addition of a spreader sticker may improve efficacy. Spray adjuvants with spreading / penetrating characteristics may improve leaf uptake and systemic concentration of active ingredient.	
Reapply at 14-28 days interval, if needed.	
Use Restrictions	
Do not apply more than 5 fl oz of product (150 mL, 0.08 lb active ingredient) per acre in a single application.	
Do not apply more than 25 fl oz of product (750 mL, 0.4 lb active ingredient) of product per acre per year.	

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577.

For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937).

As with any crop-protection product, always read and follow label directions.

For additional information call toll-free 1-866-99BAYER (1-866-992-2937).

**Section 24(c) Registrant
Bayer Environmental Science
A Division of CropScience LP
5000 GentreGreen Way, Suite 400
Cary, NC 27513**