Environmental Science US, LLC 5000 CentreGreen Way, Suite 400 Cary, NC 27513

FIFRA Section 2(ee) Recommendation

PROTHIOCONAZOLE

Densicor® Fungicide EPA Reg. No. 101563-210

For Control of Cream Leaf Blight and Ink Spot on Turfgrass

For Distribution and Use Only in the States of: ALABAMA, ARIZONA, ARKANSAS, COLORADO, DELEWARE, FLORIDA, GEORGIA, HAWAII, IOWA, IDAHO, ILLINOIS, INDIANA, KANSAS, KENTUCKY, LOUISIANA, MAINE, MARYLAND, MASSACHUSETTES, MICHIGAN, MINNISOTA, MISSISSIPPI, MISSOURI, MONTANA, NEBRASKA, NEW HAMPSHIRE, NEW JERSEY, NEVADA, NORTH CAROLINA, NORTH DAKOTA, OHIO, OKLAHOMA, OREGON, PENNSYLVANIA, RHODE ISLAND, SOUTH CAROLINA, SOUTH DAKOTA, TENNESSEE, TEXAS, UTAH, VERMONT, VIRGINIA, WASHINGTON, WISCONSIN, WEST VIRGINIA, AND WYOMING.

FIFRA Section 2(ee) Recommendation: This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA. This product bulletin should be in the possession of the user at the time of application of the product. All applicable directions, restrictions and precautions on the EPA registered label are to be followed.

DIRECTIONS FOR USE: It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This label must be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

TURF USE DIRECTIONS: DENSICOR controls a range of turf diseases. Refer to the table below for disease, rate structure, and specific use information. Apply DENSICOR prior to turf disease development or in the early stages of turf disease development. See 'Application Directions' for further information on equipment, nozzles, spray volumes and mixing instructions.

Disease	Application Rate (fl oz Product /Acre) ¹	Interval between Applications (days) ²	Use Information
Cream Leaf Blight (Limonomyces roseipellis)	8.5	14-28	Begin preventative applications when conditions are conducive to disease development.
Ink Spot (<i>Curvularia malina</i>)			

¹8.5 fl oz product/A equates to 0.196 fl oz product/1,000 ft2 or 0.267 lbs a.i./A

RATE RESTRICTIONS

- 1) Maximum Single Rate: Do not exceed 8.5 fl oz product/A per application (0.196 fl oz product/1,000 ft²), equivalent to 0.267 lbs a.i./A
- 2) Yearly Maximum Rate: Do not exceed 25.6 fl oz product/A per year (0.588 fl oz product/1,000 ft²/year), equivalent to 0.801 lbs a.i./A/year
- Do not exceed 3 applications per year using the maximum single application rate.

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577.

For PRODUCT USE Information Call 1-800-331-2867.

As with any crop-protection product, always read and follow label directions. For additional information call toll-free 1-800-331-2867).

²Use the shorter specified application interval when prolonged favorable turf disease conditions exist.