



FIFRA 2(ee) Recommendation

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE(S) OF: ALABAMA, ARIZONA, ARKANSAS, CALIFORNIA, FLORIDA, GEORGIA, HAWAII, KENTUCKY, LOUISIANA, MARYLAND, MISSISSIPPI, MISSOURI, NEW MEXICO, NORTH CAROLINA, OKLAHOMA, SOUTH CAROLINA, TENNESSEE, TEXAS, and VIRGINIA.

This recommendation is valid until December 31, 2027

Mirage[®] Stressgard[®] Fungicide

EPA Reg. No. 432-1529

Reduced Application Rate of Mirage[®] Stressgard[®] Only when Tank-Mixed with Castlon[™] Fungicide

FIFRA Section 2(ee) Recommendation: This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA. This product bulletin should be in the possession of the user at the time of application of the product. All applicable directions, restrictions and precautions on the EPA registered label are to be followed.

DIRECTIONS FOR USE:

It is a violation of federal law to use this product in a manner inconsistent with its label. This label must be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

RESTRICTIONS:

Do not use on home lawns and turf sites associated with apartment buildings, daycare centers, playgrounds, playfields, recreational parks, athletic fields, athletic fields located on or next to schools, campgrounds, churches, and theme parks. Do not use on turf grown for sale or commercial use as sod. Not for residential use.

TURF-SPECIFIC USE DIRECTIONS¹

MIRAGE STRESSGARD controls the diseases listed below

Disease	Application Rate (fl oz Product /1000 sq ft)	Interval between Applications (days)	Use Information
Fairy Ring (caused by Basidiomycete fungi)	0.6 fl oz – 2.0 fl oz	28	Water-in the treatment with the next irrigation cycle or within 10 hours to the depth at which fairy ring is present. Do not apply to overseeded bermudagrass during spring transition. For preventative applications where light disease pressure is anticipated, use the lower rate. Under high disease pressure or if there is prior history or early curative application, use the higher rate.
Large Patch (Zoysia Patch) (<i>Rhizoctonia solani</i>)	0.6 fl oz – 2.0 fl oz	28	For preventative applications where light disease pressure is anticipated, use the lower rate and longer interval. Under high disease pressure or if there is prior history or for early curative application, use the higher rate and shorter interval.
Leaf and Sheath Blight, Mini Ring (<i>Rhizoctonia zeae</i>)	0.6 fl oz – 2.0 fl oz	21 - 28	Apply preventively or curatively when conditions are favorable for disease development.
Spring Dead Spot (<i>Ophiosphaerella korrae</i> , <i>O. herpotricha</i> , <i>Leptosphaeria korrea</i> , <i>L. Namari</i>)	0.6 fl oz – 2.0 fl oz	28	Lightly water-in applications to move fungicide into crown area and root zone.
Take-all Root Rot, Bermudagrass Decline, Warm Season Turfgrass Decline (<i>Gaeumannomyces graminis</i> var. <i>graminis</i>)	0.6 fl oz – 2.0 fl oz	28	Apply fungicide in adequate water volume or water-in applications to upper root zone. Apply fungicide prior to periods of stress, including hot, humid conditions or extended wet weather.

¹Do not exceed 6.5 fl oz of Mirage Stressgard per 1,000 sq ft per year.

For all other use directions, restrictions, and precautions please refer to the Mirage Stressgard EPA-registered label.

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577.

For PRODUCT USE Information Call 1-800-331-2867.

As with any crop-protection product, always read and follow label directions.

For additional information call toll-free 1-800-331-2867).

**Bayer Environmental Science,
A Division of Bayer CropScience LP
5000 CentreGreen Way, Suite 400
Cary, NC 27513**