


MAXFORCE® Ant and Cockroach Bait Granules

REG. NO. L5658 Act No. 36 of 1947
Before using this product read the label carefully

READY-TO-USE GRANULAR BAIT FOR THE CONTROL OF ANTS (INCLUDING BROWN HOUSE ANT AND BLACK SUGAR ANT) AND COCKROACH OUTDOORS AROUND DOMESTIC PREMISES AND BUILDINGS, ON LAWNS, TURFGRASS AND OTHER AREAS SUCH AS PARKS AND GOLF COURSES.

	<p>Signal word: Danger</p> <p>Hazard statements Causes damage to organs through prolonged or repeated exposure. Very toxic to aquatic life with long lasting effects.</p> <p>Precautionary statements Do not breathe dust Wear protective gloves/ protective clothing/ eye protection/ face protection</p>
---	--

ACTIVE INGREDIENT:

Hydramethylnon 10 g/kg

NETTO CONTENTS: g

REGISTRATION HOLDER:

2022 Environmental Science ZA (Pty) Ltd.
Reg. No. / Nr. 2022/228577/07
P.O. Box 143
ISANDO 1600
SOUTH AFRICA / SUID-AFRIKA
TEL: (011) 921 5911

For any product enquiry, contact 2022 Environmental Science ZA (Pty) Ltd. on 011 921 5911

For poisoning emergencies only, Griffon Poison Information Centre may also be contacted - 082 446 8946. National Poison Centre - + 27 (0861) 555 777
Transport / Spill emergency no: (Chemtrec) +27 80 098 3611 (2022 Environmental Science ZA Pty Ltd)

Batch no:
Date manufactured:
Shelf life: 2 years after date of manufacture

MAXFORCE® is a registered trademark of the Bayer Group.

UN No. 3077

WARNINGS:**Hazard statements**

May damage fertility.

To avoid risks to human health and the environment, comply with the instructions for use.

Contains 2-tert-Butylhydroquinone. May produce an allergic reaction.

Harmful if swallowed.

May irritate skin and eyes.

Toxic to fish. [Prevent contamination of fishponds, streams, or rivers.](#)Keep out of reach of children, uninformed persons ([bystanders](#)) and animals.

Store in a cool, dry place, away from food, feedstuffs and other insecticides.

Keep container closed when not in use.

DISCLAIMER:

ALTHOUGH THIS PRODUCT HAS BEEN EXTENSIVELY TESTED UNDER A LARGE VARIETY OF CONDITIONS, FAILURE THEREOF MAY ENSUE AS A RESULT OF A WIDE RANGE OF REASONS. IF THIS IS SUSPECTED, KINDLY NOTIFY THE REGISTRATION HOLDER.

PRECAUTIONS:**Precautionary statements**

IF exposed or concerned: Call a POISON CENTER/ doctor/ physician.

Dispose of contents/container in accordance with local regulation.

[Avoid contamination to waterways. Apply the product to specific cracks and crevices in paving or concrete and avoid general broadcast applications. Avoid making applications if it is likely to rain within 24 hours of application.](#)

Avoid contact with skin or eyes.

Wear gloves and wash hands thoroughly after use.

Avoid contamination with other insecticides.

Destroy empty container and do not re-use for any other purpose.

RELEVANT SUBSTANCES

Name	CAS-No. / EC-No. / REACH Reg. No.	Conc. [%]
Hydramethylnon	67485-29-4	1,0
Oleic acid	112-80-1	>= 1,0 – <= 5,0
2-tert-Butylhydroquinone	1948-33-0	>= 0,1 – <= 1,0

DIRECTIONS FOR USE: USE ONLY AS DIRECTED.

For outdoor use only.

The delayed action of the active ingredient allows foraging ants to take the bait back to the nest so that the entire colony, including the queen, is destroyed. When properly used, this bait will control many common structure-invading ants before they enter a building. Do not apply other insecticides e.g. aerosols, sprays, etc. on top of or in close proximity to **Maxforce® Ant and Cockroach Bait Granules**, as it will reduce the effectiveness. Reduced activity and dead ants around the entrance of the nest should be noted around 7

days after baiting. [Follow product application instructions on the label and excessive application of the product is discouraged. The number of applications in residential areas should not exceed 4 times per year.](#)

1. Perimeter treatment

Lightly sprinkle granules at the rate of 6 g/m² in a band 0,5 m wide around the perimeter of buildings where ants are active. Treat ant trails, cracks and crevices where ants may enter structures.

2. Broadcast application

Broadcast granules uniformly on lawns and other areas at the rate of 20 g/100 m². Bait will remain effective, even after exposure to rain and sprinklers, for up to 7 days in direct sunlight.

3. Ant nests

Sprinkle approximately 30 g bait around active ant nests. Do not apply bait directly in or on the mound. If the mound is disturbed, ants will become agitated and may not feed on the bait.

4. Cockroaches

Indoor application

Lightly sprinkle granules at the rate of 0.2 g/m² into cracks and crevices where the cockroaches are active. Remove any loose or excess granules. Do not use in areas accessible by children, uninformed persons and animals.

MAXFORCE® is a registered trademark of the Bayer Group.

© 2022 Environmental Science ZA (Pty) Ltd. All unauthorized reproduction of this label is forbidden.

www.za.envu.com